

New Record of *Scolecenchelys fuscogularis* (Anguilliformes: Ophichthidae) Leptocephali from Korea, as Revealed by Morphological and Molecular Analyses

Hwan-Sung Ji¹, Jin-Koo Kim^{2*}, Soo Jeong Lee², Seishi Kimura³ and Yusuke Hibino³

¹Fisheries Resources Management Division, National Fisheries Research and Development Institute, Busan 619-705, Korea

²Department of Marine Biology, Pukyong National University, Busan 608-737, Korea

³Fisheries Research Laboratory, Mie University, 4190-172, Wagu, Shima-cho, Shima, Mie 517-0703, Japan

Abstract

Three leptocephali (22.2, 22.7, 56.0 mm in total length) collected from the East/Japan Sea were identified by morphological and genetic analyses as belonging to the genus *Scolecenchelys* (Anguilliformes, Ophichthidae). Morphologically, the specimens were characterized by 148-158 myomeres, 10 gut swellings, dorsal fin origin above middle of the body, and 6 postanal melanophores between the anus and the caudal margin. An analysis of an 849-base pair 12S rRNA sequence of mitochondrial DNA showed that sequences are concordant with those of adult *Scolecenchelys fuscogularis* (genetic distance = 0.001). Furthermore total number of myomeres is consistent with the total number of vertebrae in adult *S. fuscogularis*. This study provides the first description of the morphological characteristics of *S. fuscogularis* leptocephali and their variations with size. The Korean name of *S. fuscogularis* is "Ga-neun-mul-baem", established by Ji et al. (2012).

Key words: *Scolecenchelys fuscogularis*, New record, Ophichthidae, Leptocephalus, mtDNA 12S rRNA

Introduction

The genus *Scolecenchelys* Ogilby, 1897, of the subfamily Myrophinae, family Ophichthidae, order Anguilliformes, is distributed in tropical and temperate areas of the Indo-Pacific (Hibino et al., 2013a). The genus includes 21 species (Hibino et al., 2013a; Eschmeyer, 2014), 7 of which are known in Japan (Hataoka, 2013; Hibino et al., 2013a, b). However, only one species, *Scolecenchelys aoki* (Jordan and Snyder, 1901) (known from adults and leptocephali) has been recorded from Korea (Ji et al., 2012; Ji et al., 2013). Species of *Scolecenchelys* in adults are defined by the following combination of morphological characters: absence of pectoral fin; center of the eye located behind the mid-jaw; presence of three preopercular sensory pores, and two infraorbital pores located between the anterior and posterior nostrils; and teeth on upper jaw and

vomer conical, pointed, and arranged in single to triple rows (Castle and McCosker, 1999; McCosker et al., 2012). Castle and McCosker (1999) provided a taxonomic review of *Scolecenchelys*, and several new species of the genus have been recently described by McCosker (2006), McCosker et al. (2012) and Hibino et al. (2013a).

During November 2009 and October 2010, three leptocephali collected from the East/Japan Sea, and were identified as *Scolecenchelys fuscogularis* (Hibino, Kai and Kimura, 2013a) based on a combination of morphological and molecular traits. Our study provides the first report of this species in Korean waters. We here present a detailed morphological description of size-related variations in the leptocephali of *S. fuscogularis*.

 © 2015 The Korean Society of Fisheries and Aquatic Science

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Received 29 October 2014; Revised 20 November 2014

Accepted 20 November 2014

*Corresponding Author

E-mail: taengko@hanmail.net




Fig. 1. Leptocephalus of *Scolecenchelys fuscogularis*, PKU 5999, 22.2 mm TL (A); PKUI 205, 56.0 mm TL (B). Scale bars= 1.5 mm. DO: dorsal fin origin.

Materials and Methods

Material examined : PKU 5999-6000, 2 specimens, 22.2 mm and 27.2 mm in total length (TL), southern part of the East/Japan Sea (35°40'43"N, 129°44'47"E), October 2010, RN80 net, collected by JK Kim and HS Ji. PKUI 205, 1 specimen, 56.0 mm TL, southern part of the East/Japan Sea (34°54'47", 129°33'14"E), 22 November 2009, RN80 net, collected by JK Kim and HS Ji.

The specimens were immediately preserved in 99% ethanol on shipboard. Counts and measurements followed those of Fahay and Obenchain (1978) and Tabeta and Mochioka (1988). Each body part was measured to the nearest 0.01 mm using digital vernier calipers, with measurements performed under zoom stereomicroscope (Olympus SZX-16, Japan). The description of tooth sequences followed Castle (1984). The specimens collected in the study are deposited in the Ichthyoplankton Collection of Pukyong National University (PKUI), Korea.

Genomic DNA was extracted from the right eyeball of the *S. fuscogularis* leptocephali (22.2–56.0 mm TL) using forceps and Chelex 100 resin (Bio-Rad, USA). We also analyzed DNA of the adult *S. fuscogularis* using muscle tissue obtained from the holotype of the species, FAKU (Faculty of Agriculture, Kyoto University, Japan) 132857. Later, mitochondrial 12S rRNA sequences were amplified and sequenced according to the method of Ji and Kim (2010). The DNA sequences were aligned and edited using Clustal W (Thompson et al., 1994) and BioEdit version 7.0.0. The molecular results were compared with data from five adult ophichthids (*S. aoki*, *Muraenichthys gymnopterus*, *Echelus uropterus*, *Pisodonophis sangjuensis*, and *Ophichthus asakusae*) and one outgroup (*Conger myriaster*). Genetic divergences were calculated using the Kimura two-parameter model (Kimura, 1980) and MEGA 6 (Tamura et al., 2013); a neighbor-joining (NJ) tree was constructed using the Kimura two-parameter method and 10,000 bootstrap replications in MEGA 6.

Results and Discussion

Scolecenchelys fuscogularis Hibino, Kai and Kimura, 2013

(Korean name: Ga-neun-mul-baem; Fig. 1, Table 1)

Scolecenchelys fuscogularis Hibino, Kai and Kimura, 2013: 43 (type locality: the Sea of Japan off the coast of Kyoto Prefecture, Japan).

Scolecenchelys sp.: Ji et al., 2012: 415 (Korea).

Identification

Morphological analysis

The present specimens have following characters: first and second liver lobe separated; strongly developed pterygiophores; and dorsal fin origin slightly above and behind the midpoint of the body. From these characters, the present leptocephali can be decided as a member of the subfamily Myrophinae (Castle, 1984). The present specimens also have following characters: 148-158 total myomeres; 10 gut swellings; dorsal fin origin above middle of the body; and 6 postanal melanophores between the anus and the caudal margin, and these characters are shared with a leptocephalus of *Scolecenchelys aoki* (Ji et al., 2012).

Mitochondrial DNA sequence analysis

The 849-base pair sequences of mtDNA 12S rRNA in the three leptocephali in this study were nearly identical to the sequences of adult *S. fuscogularis* (genetic distance, $d = 0.001$), but were slightly distinct from adult *S. aoki* ($d = 0.006$) (Table 2). The neighbor joining tree showed that the three leptocephali were clustered with adult *S. fuscogularis*, being strongly supported by a 98% bootstrap value (Fig. 2).




Fig. 2. Neighbor-joining (NJ) tree based on partial mtDNA 12S rRNA sequences showing the relationships among the three *Scolecenchelys fuscogularis* leptocephali and 6 adult ophichthids with one outgroup *Conger myriaster*. The NJ tree was constructed using the Kimura-2-parameter distance model. 10,000 replications of bootstrap were conducted. Bar indicates genetic distance of 0.02.

Table 1. Comparisons of measurements and counts for *Scolecenchelys fuscogularis* and *Scolecenchelys aoki*

Species	<i>S. fuscogularis</i> (present study)	<i>S. fuscogularis</i> (present study)	<i>S. aoki</i> (Ji et al., 2012)	<i>S. fuscogularis</i> (Hibino et al., 2013a)
Development	Caudal fin absent	Caudal fin development	Caudal fin development	adult
The number of specimens	2	1	1	2
Total length (mm, TL)	22.2-27.2	56.0	59.2	187.0-265.0
In % of total length				
Head length	8.1-9.5	6.8	7.4	-
Predorsal length	-	59.5	53.5	-
Preanal length	65.4-68.0	59.9	57.7	-
Body depth	7.2-7.7	8.6	10.1	-
In % of Head length				
Eye diameters	18.2-19.0	13.2	11.4	-
Snout length	42.9-45.5	34.2	25.0	-
Upper jaw length	56.8-57.8	52.6	55.2	-
Counts				
Total myomeres	148-158	150	135	146-149 ^I
Predorsal myomeres	-	74	64	72 ^{II}
Preanal myomeres	73-74	71	69	63 ^{III}
1st vertical blood vessel	9-11	10	16	-
Last vertical blood vessel	69	69	63	-
1st gut swelling	10-12	11	12	-
Last gut swelling	71-72	69	67	-
Postanal pigment	6-8	6	-	-
Number of nostril	2	2	2	-
Number of gut swelling	10	10	10	-
Dentition formular	$\frac{1 + II - III + 1 - 2}{1 + III + 1 - 2}$	$\frac{1 + IV + 4}{1 + IV + 3}$	$\frac{1 + IV + 4}{1 + IV + 3}$	-

Superscript indicate number of total vertebrae (I), predorsal vertebrae (II) and preanal vertebrae (III).

Table 2. Genetic distance among *Scolecenchelys fuscogularis* leptocephali, 6 adult ophichthid species and 1 outgroup

	(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)
<i>Scolecenchelys fuscogularis</i> leptocephalus (1)									
<i>Scolecenchelys fuscogularis</i> leptocephalus (2)	0.000								
<i>Scolecenchelys fuscogularis</i> leptocephalus (3)	0.000	0.000							
<i>Scolecenchelys fuscogularis</i> adult (4)	0.001	0.001	0.001						
<i>Scolecenchelys aoki</i> adult (5)	0.006	0.006	0.006	0.006					
<i>Muraenichthys gymnopterus</i> adult (6)	0.109	0.109	0.109	0.109	0.108				
<i>Echelus uropterus</i> adult (7)	0.166	0.166	0.166	0.166	0.161	0.195			
<i>Pisodonophis sangjuensis</i> adult (8)	0.184	0.184	0.184	0.184	0.182	0.223	0.125		
<i>Ophichthus asakusae</i> adult (9)	0.187	0.187	0.187	0.187	0.185	0.217	0.084	0.097	
<i>Conger myriaster</i> adult (10)	0.206	0.206	0.206	0.206	0.204	0.198	0.214	0.217	0.228

Description of *leptocephalus* of *Scolecenchelys fuscogularis*

Precaudal fin developmental stage (22.2 mm and 27.2 mm TL): Head relatively small; tip of snout highly acute. Fang-like teeth on both jaws. Two nostrils present in front of eyes. Anterior tip of lower jaw more acute than that of upper jaw. Caudal region short. Head length, 8.1-9.5% TL; body depth, 7.2-7.7% TL; preanal length, 65.4-78.0% TL (Table 1). Punctate and branched melanophores present on dorsal surfaces of all 10 gut swellings (Fig. 1A); six developed melanophores present just below the lateral midline of the caudal region and extending to the caudal terminus (Fig. 1A).

Caudal fin developmental stage (56.0 mm TL): Number of teeth increasing from the preceding stage. Unlike in the preceding stage, both jaws coincident (Fig. 1B). Head length (6.8% TL) and preanal length (59.9% TL) becoming smaller than those of the preceding stage, whereas body depth (8.6 % TL) larger (Table 1). All fins developed; dorsal fin origin located above 74th myomere, anal, caudal, and pectoral fin rays developed. Unlike in the preceding stage, the 6 melanophores on the caudal region fading (Fig. 1B).

Distribution

East/Japan Sea (Ji et al., 2012; Hibino et al., 2013a; present study), and off the Pacific coast of Japan (Hibino et al., 2013a).

Remarks

The total number of myomeres in the present *S. fuscogularis* leptocephali (148-158) is approximate to the range of the number of vertebrae in adult *S. fuscogularis* (146-149) (Hibino et al., 2013a), and the number of predorsal myomeres in the present leptocephali (74) is also consistent with number of predorsal vertebrae in adult *S. fuscogularis* (72) (Table 1; Hibino et al., 2013). Although *S. fuscogularis* are most similar to *Scolecenchelys australis* (Macleay, 1881) and *Scolecenchelys chilensis* (McCosker, 1970) in the number of vertebrae (146-149 in *S. fuscogularis* vs. 148-152 in *S. australis* vs. 146-159 in *S. chilensis*) (Hibino et al., 2013a), their distributional areas are different (northeast Asia for *S. fuscogularis* vs. the south eastern Pacific for *S. australis* vs. the south western Pacific for *S. chilensis*) (McCosker, 1970; 1977; Hibino et al., 2013a). *Scolecenchelys fuscogularis* leptocephali differ from *S. aoki* leptocephali in the number of myomeres (148-158 in *S. fuscogularis* vs. 135 in *S. aoki*), and the number of preanal myomeres (71-74 in *S. fuscogularis* vs. 69 in *S. aoki*) (Table 1; Ji et al., 2012). We adopted the Korean name for *S. fuscogularis*, “Ga-neun-mul-baem”, following Ji et al. (2012).

Acknowledgments

We are deeply indebted to Dr. Y. Kai (FAKU) for his donating the tissue sample of the holotype *Scolecenchelys fuscogularis*. This research was supported by the Marine Biodiversity Institute of Korea (MABIK).

References

- Castle PHJ. 1984. Notacanthiformes and Anguilliformes: Development. In: Ontogeny and systematics of fishes. Moser HG, Richards WJ, Cohen DM, Fahay MP, Kendall AW and Richardson SL, eds. Allen Press INC., Lawrence, USA, 62-69.
- Castle PHJ and McCosker JE. 1999. A new genus and two new species of Myrophinae worm-eels, with comments on *Muraenichthys* and *Scolecenchelys* (Anguilliformes: Ophichthidae). Rec Aust Mus 51, 113-122.
- Eschmeyer WN. ed. 2014. Catalog of Fishes Electronic version. available from: <http://research.calacademy.org/ichthyology/catalog/fishcatmain.asp>.
- Fahay MP and Obenchain CL. 1978. Leptocephali of the ophichthid genera *Ahlia*, *Myrophis*, *Ophichthus*, *Pisodonophis*, *Callechelys*, *Letharchus*, and *Apterichtus* on the atlantic continental shelf of the United States. Bull Mar Sci 28, 442-486.
- Hatooka K. 2013. Ophichthidae. In: Fishes of Japan with Pictorial Keys to the Species. Third ed. Nakabo T, ed. Tokai University Press, Kanagawa, Japan, 266-277, 1794-1802.
- Hibino Y, Kai Y and Kimura S. 2013a. *Scolecenchelys fuscogularis* (Anguilliformes: Ophichthidae, Myrophinae), a new worm eel from Japan. Ichthyol Res 60, 43-47.
- Hibino Y, Kimura S and Hatooka K. 2013b. First records of a worm eel (Anguilliformes: Ophichthidae, Myrophinae), *Scolecenchelys laticaudata* from Kikai Island, Kagoshima, Japan. Jpn J Ichthyol 60, 35-41.
- Ji HS and Kim JK. 2010. Molecular and morphological identification of ophichthid leptocephali from the south sea of Korea. Korean J Ichthyol 22, 279-284.
- Ji HS, Lee HW, Hong BK and Kim JK. 2012. Four newly reported ophichthid leptocephali species revealed by mitochondrial 12S rDNA, with implications of their occurrence in Korea. Anim Cells Syst 16, 415-424.
- Ji HS, Kim MJ and Kim JK. 2013. First description of the adult *Scolecenchelys aoki* (Ophichthidae, Anguilliformes) collected from the South Sea of Korea, with comments on the synonym *Scolecenchelys borealis*. Kor J Fish Aquat Sci 46, 589-593.
- Jordan DS and Snyder JO. 1901. A review of the apodal fishes or eels of Japan, with descriptions of nineteen new species. Proc US Natl Mus 23, 837-890.
- Kimura M. 1980. A simple method for estimating evolutionary rates of base substitutions through comparative studies of nucleotide sequences. J Mol Evol 16, 111-120.
- Macleay W. 1881. A descriptive catalogue of Australian fishes. Part IV. Proc Linn Soc NSW 6, 202-387.

- McCosker, JE. 1970. A review of the eel genera *Leptenchelys* and *Muraenichthys*, with the description of a new genus, *Schismorhynchus*, and a new species, *Muraenichthys chilensis*. Pac Sci 24, 506-516.
- McCosker JE. 1977. The osteology, classification, and relationships of the eel family Ophichthidae. Proc Calif Acad Sci Ser 4 41, 1-123.
- McCosker JE. 2006. A new deepwater species of worm-eel, *Scolecenchelys castlei* (Anguilliformes: Ophichthidae), from New Zealand and Australia, with comments on *S. breviceps* and *S. macropoda*. J Royal Soc 36, 17-26.
- McCosker JE, Ide S and Endo H. 2012. Three new species of ophichthid eels (Anguilliformes: Ophichthidae) from Japan. Bull Natl Mus Nat Sci (Ser A) Suppl 6, 1-16.
- Ogilby JD. 1897. Some new genera and species of fishes. Proc Linn Soc NSW 22, 245-251.
- Tabeta O and Mochioka N. 1988. Family Ophichthidae. In: An Atlas of the Early Stage Fishes in Japan. Okiyama M, ed. Tokai University Press, Tokyo, Japan, 58-62.
- Tamura K, Stecher G, Peterson D, Filipski A, Kumar S. 2013. MEGA6: molecular evolutionary genetics analysis version 6.0 Mol Bio Evol 30, 2725-2729.
- Thompson JD, Higgins DG and Gibson TJ. 1994. CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position specific gap penalties and weight matrix choice. Nucl Acids Res 22, 4673-4680.